

MYCOTAXON

DOI: 10.5248/113.263

Volume 113, pp. 263–267

July–September 2010

A new *Trametes* species from Southwest China

HAI-JIAO LI & BAO-KAI CUI*

* *baokaicui@yahoo.com.cn**Institute of Microbiology, P.O. Box 61, Beijing Forestry University
Beijing 100083, China*

Abstract — A new polypore, *Trametes cystidiolophora* sp. nov., found in Yunnan Province, southwest China, is described and illustrated. The new species is characterized by its pale grayish brown to pale cinnamon-buff pileus with distinctly concentric zones and radial veins, uneven pore surface, cylindrical to more or less allantoid basidiospores ($6.6\text{--}9.2 \times 2.4\text{--}3 \mu\text{m}$), abundant cystidioles present in the hymenium, and skeletal hyphae that become swollen in KOH.

Key words — *Polyporaceae*, taxonomy, wood-rotting fungi

Introduction

The genus *Trametes* Fr. is characterized by having pileate basidiocarps, a trimitic hyphal system with clamp connections on generative hyphae, hyaline and thin-walled basidiospores that are negative in Melzer's reagent, and causing white rot (Gilbertson & Ryvarden 1986, Ryvarden & Gilbertson 1994, Lindblad & Ryvarden 1999, Núñez & Ryvarden 2001). About 50 species in the genus have been reported in the world (Kirk et al. 2008), including 23 species previously recorded from China (Zhao & Zhang 1991, Teng 1996, Zhao 1998, Dai et al. 2007, Dai 2009, Dai & Yuan 2010).

During the study on wood-decaying fungi from Gaoligongshan Nature Reserve, Yunnan Province, southwest China, a species of *Trametes* was found that could not be identified to any known species. It is described in the present paper as *Trametes cystidiolophora*.

Materials and methods

The studied specimens are deposited in herbaria as cited below. The microscopic procedure follows Dai & Penttilä (2006). Sections were studied at

* Corresponding author

magnification up to $\times 1000$ using a Nikon Eclipse E 80i microscope and phase contrast illumination. Drawings were made with the aid of a drawing tube. To present spore size variation, the 5% of measurements excluded from each end of the range are given in parentheses. Abbreviations include IKI = Melzer's reagent, IKI- = negative in Melzer's reagent, KOH = 5% potassium hydroxide, CB = Cotton Blue, CB- = acyanophilous, L = mean spore length (arithmetic average of all spores), W = mean spore width (arithmetic average of all spores), Q = variation in the L/W ratios between the specimens studied, and n = number of spores measured from given number of specimens. Special colour terms follow Anonymous (1969) and Petersen (1996).

Taxonomy

Trametes cystidiolophora B.K. Cui & H.J. Li, sp. nov.

FIG. 1

MYCOBANK MB 518544

Carpophorum annuum, *pileatum*, *imbricatum*; *facies pororum bubalina vel reseo-bubalina*, *pori rotundi vel angulati*, 2–3 per mm. *Systema hypharum trimiticum*, *hyphae generativae hyalinae, fibulatae*, *hyphae skeletales contexti* 2.8–6.2 μm ; *spores hyalinae, cylindricae*, IKI-, CB-, 6.6–9.2 $\times$ 2.4–3 μm .

TYPE — China. Yunnan Province, Baoshan, Gaoligongshan Nature Reserve, on dead angiosperm tree, 25.X.2009 Cui 8084 (holotype in BJFC); Cui 8087 (isotype in BJFC).

ETYMOLOGY — *cystidiolophora* (Greek): = “cystidiole-bearing”, referring to the abundant cystidioles in the hymenium.

FRUITBODY — Basidiocarps annual, pileate, usually imbricate, without odor or taste when fresh, corky and light in weight when dry. Pileus dimidiate to semicircular, projecting up to 4.2 cm, 7.3 cm wide, 7 mm thick at the base; pileal surface pale grayish brown to pale cinnamon-buff when dry, glabrous, distinctly concentrically zoned and radially veined; margin sharp, wavy or incised in rounded lobes, deflexed with age. Pore surface cream-buff to pinkish buff when dry, slightly shiny; sterile margin white to cream, up to 2.5 mm wide; pores round to angular, 2–3 per mm, dissepiments thin, entire at margin and dentate to hydroid with age. Context cream, corky, up to 3 mm thick. Tubes cream to cream-buff, corky, up to 4 mm long.

HYPHAL STRUCTURE — Hyphal system trimitic; generative hyphae with clamp connections; skeleto-binding hyphae dominant, thick-walled to subsolid, IKI-, CB-, become swollen in KOH.

CONTEXT — Generative hyphae infrequent, hyaline, thin-walled, moderately branched, 2–3.7 μm in diam; skeletal hyphae dominant, hyaline, slightly thick-walled to subsolid, frequently branched, and the slightly thick-walled skeletal hyphae often collapsed, interwoven, 2.8–6.2 μm in diam; binding hyphae hyaline, thick-walled to almost solid, frequently branched, interwoven, 1.7–3 μm .




FIG. 1. Microscopic structures of *Trametes cystidiolophora* (drawn from the holotype).
a: Basidiospores. b: Basidia and basidioles. c: Cystidioles.
d: Hyphae from tube trama. e: Hyphae from context.

TUBES — Generative hyphae infrequent, hyaline, thin-walled, frequently branched, 1.7–3 µm in diam; skeletal hyphae dominant, hyaline, thick-walled to subsolid, occasionally branched, interwoven, 2.3–5 µm; binding hyphae hyaline, flexuous, thick-walled to almost solid, frequently branched, interwoven, 1.6–3.1 µm. Cystidia absent, cystidioles abundant in the hymenium, fusoid, hyaline, mostly thin-walled, occasionally slightly thick-walled, some with one or two septa, 16–24 × 4–6 µm; basidia clavate, with four sterigmata and a basal clamp connection, 16–18.2 × 5–7.8 µm; basidioles in shape similar to basidia, but slightly smaller.

SPORES — Basidiospores cylindrical, occasionally slightly curved to more or less allantoid hyaline, thin-walled, smooth, IKI–, CB–, (6–)6.6–9.2(–10) × (2.2–)2.4–3(–3.3) µm, L = 8.1 µm, W = 2.79 µm, Q = 2.78–3.04 (n=60/2).

TYPE OF ROT — White rot.

REMARKS — *Trametes cystidiolophora* is characterized by its pale grayish brown to pale cinnamon-buff pileal surface with distinctly concentric zones and radial veins, uneven pore surface, cylindrical to more or less allantoid basidiospores (6.6–9.2 × 2.4–3 µm), and abundant cystidioles present in the hymenium. Moreover, its skeleto-binding hyphae become swollen in KOH.

Trametes cystidiolophora may be confused with *Fomitopsis palustris* (Berk. & M.A. Curtis) Gilb. & Ryvarden, which produces similar basidiospores, but the two species can be separated by the rot type. *Trametes cystidiolophora* causes a white rot, while *Fomitopsis palustris* causes a brown rot.

Trametes maxima (Mont.) A. David & Rajchenb. is similar to *T. cystidiolophora* by sharing similar uneven pore surface and pore size (2–3 per mm), but *T. maxima* differs in its tomentose to hirsute pileal surface and smaller basidiospores (4.5–5.5 × 2–2.5 µm, Gilbertson & Ryvarden 1986).

Trametes cotonea (Pat. & Har.) Ryvarden, which has similar basidiospores (7–10 × 2.5–3.5 µm) as *T. cystidiolophora*, is usually effused-reflexed, paper thin, and flexible and its pores are smaller (3–4 per mm, Ryvarden & Johansen 1980).

Trametes glabrata (Lloyd) Ryvarden, which shares with *T. cystidiolophora* a glabrous pileus, uneven pore surface, and similar pore size. However, *T. glabrata* has distinctly smaller basidiospores (4–5 × 1–1.5 µm, Ryvarden 1992).

Cystidioles are also present in several other reported species in *Trametes*, such as *T. gibbosa*, *T. hirsuta*, *T. ljubarskyi*, and *T. pubescens* (Gilbertson & Ryvarden 1986, Núñez & Ryvarden 2001), but the cystidioles in all these species are infrequent and without septa. The fact that its cystidioles are abundant in the hymenium and that some are septate make *T. cystidiolophora* unique in *Trametes*.

Acknowledgements

We express our gratitude to Drs. Yu-Cheng Dai (Shenyang, China) and Wjacheslav A. Spirin (St.Petersburg, Russia) who reviewed the manuscript. The research was financed by the National Natural Science Foundation of China (Project No. 30900006), the Fundamental Research Funds for the Central Universities (Project No. BLYX200912) and the Ministry of Science and Technology of China (Project No. 2008BADB0B03).

Literature cited

- Anonymous. 1969. Flora of British fungi. Colour identification chart. Her Majesty's Stationery Office, London. 1 pp.
- Dai YC. 2009. A checklist of polypores in China. *Mycosystema* 28: 315–327.
- Dai YC, Cui BK, Huang MY. 2007. Polypores from eastern Inner Mongolia, northeastern China. *Nova Hedwigia* 84: 513–520. doi:10.1127/0029-5035/2007/0084-0513
- Dai YC, Penttilä R. 2006. Polypore diversity of Fenglin Nature Reserve, northeastern China. *Annales Bot. Fennici* 43: 81–96.
- Dai YC, Yuan HS. 2010. Type studies on polypores described by J.D. Zhao. *Annales Bot. Fennici* 47: 113–117.
- Gilbertson RL, Ryvarden L. 1986. North American polypores 2. *Megasporoporia – Wrightoporia*. Fungiflora, Oslo. 434–885 pp.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. Dictionary of the Fungi 10th. CAB International, Wallingford, Oxon. 771 pp.
- Lindblad I, Ryvarden L. 1999. Studies in neotropical polypores 3. New and interesting *Basidiomycetes* (*Poriales*) from Costa Rica. *Mycotaxon* 71: 335–359.
- Núñez M, Ryvarden L. 2001. East Asian polypores 2. *Polyporaceae* s. lato. *Syn. Fungorum* 14: 165–522.
- Petersen JH. 1996. Farvekort. The Danish Mycological Society's colour-chart. Foreningen til Svampekundskabens Fremme, Greve. 6 pp.
- Ryvarden L. 1992. Type studies in the *Polyporaceae* 23. Species described by C.G. Lloyd in *Lenzites*, *Polystictus*, *Poria* and *Trametes*. *Mycotaxon* 44: 127–136.
- Ryvarden L, Gilbertson RL. 1994. European polypores 2. *Syn. Fungorum* 7: 394–743.
- Ryvarden L, Johansen I. 1980. A preliminary polypore flora of East Africa. Fungiflora, Oslo. 636 pp.
- Teng SC. 1996. Fungi of China. Mycotaxon, Ltd, Ithaca, New York. 586 pp.
- Zhao JD. 1998. Flora Fungorum Sinicorum vol. 3 *Polyporaceae*. Science Press, Beijing. 456 pp.
- Zhao JD, Zhang XQ. 1991. Four new species of *Polyporaceae* in China. *Acta Mycologica Sinica* 10: 266–272.